arXiv:1512.05694v1 [math.NT] 17 Dec 2015

ON ONE TYPE OF COMPACTIFICATION OF POSITIVE INTEGERS

Milan Pasteka

Abstract. *The object of observation is a compact metric ring containing positive integers as dense subset. It is proved that this ring is isomorphic with a ring of reminder classes of polyadic integers.*

Let $\mathbb{N}$ be the set of positive integers. A mapping $||\cdot|| : \mathbb{N} \to < 0, \infty)$ will be called norm if and only if the following conditions are satisfied for $a, b \in \mathbb{N}$
$||a|| = 0 \Leftrightarrow a = 0$,
$||a + b|| \leq ||a|| + ||b||$,
$||ab|| \leq ||b||$.

There are various examples of norms on $\mathbb{N}$. One of these is polyadic norm defined in [N], [N1]. We start by a generalization of polyadic norm. Denote by $a + (m)$ the arithmetic progression with difference m which contains a. Instead of $0 + (m)$ we write only (m).

A subset $A \subset \mathbb{N}$ we call *closed to divisibility* or shortly CD - set if and only if
$1 \in A$,
$m \in A, d|m \Rightarrow d \in A$,
$m_1, m_2 \in A \Rightarrow [m_1, m_2] \in A$, for $d, m, m_1, m_2 \in \mathbb{N}$.

Suppose that A is infinite CD - set and $\{B_n\}$ is such sequence elements of A that for every $d \in A$ there exists n_0 that $d|B_n$ for $n > n_0$. It is easy to see that the mapping

$$||a||_A = \sum_{n=1}^{\infty} \frac{h_n(a)}{2^n}$$

for $a \in \mathbb{N}$ where $h_n(a) = 1 - \mathcal{X}_{(B_n)}$, is a norm. This norm will be called generalized polyadic norm and the completion with respect the metric given by this norm will be called the ring of generalized polyadic integers.

If $A = \mathbb{N}$ and $B_n = n!$ we get polyadic norm and the completion will be the ring of polyadic integers. In the case $A = \{p^n; n = 0, ., 2, ..\}$ and $B_n = p^n$ for given prime p we obtain p - adic norm and the completion will be the ring of p - adic integers.

In the following text we shall assume that is given a compact metric space (Ω, ρ) containing $\mathbb{N}$ as dense subset. We suppose that the operations addition and multiplication on $\mathbb{N}$ are continuous and are extended to whole Ω to continuous operations. Thus $(\Omega, +, \cdot)$ is a topological commutative semiring.

Since Ω is compact we can suppose that there exists an increasing sequence of positive integers $\{x_n\}$ convergent to an element of Ω. Put $a_n = x_{2n} - x_n, n = 1, 2, ...$ Then $a_n \geq n$ and

$$a_n \to 0 \tag{1}$$

in the topology of Ω.

For $\beta \in \Omega$ and $b_n \to \beta$, $b_n \in \mathbb{N}$ we can consider the sequence of positive integer $\{a_{k_n} - b_n\}$, for suitable increasing sequence $\{k_n\}$, that $a_{k_n} - b_n \to \beta'$ where $\beta + \beta' = 0$. We see that $(\Omega, +)$ is a compact group.

Clearly for every $m \in \mathbb{N}$ there holds $cl(r + (m)) = r + m\Omega$ where $m\Omega$ is the principal ideal in the ring $(\Omega, +, \cdot)$ generated by m. This yields

$$\Omega = m\Omega \cup (1 + m\Omega) \cup ... \cup (m - 1 + m\Omega). \tag{2}$$

Since the divisibility by m in $\mathbb{N}$ is not necessary equivalent with divisibility by m in Ω it is not provided that the last decomposition is disjoint.

Lemma 1. Let $m \in \mathbb{N}$ is such positive integer that it is also the minimal generator of the ideal $m\Omega$. Then every positive integer is divisible by m in $\mathbb{N}$ if and only if it is divisible by m in Ω.

Proof. One implication is trivial. Suppose now that some positive integer a is divisible by m in Ω. Thus $a \in m\Omega$. Put $d = (a, m)$ - the greatest common divisor in $\mathbb{N}$. Then for $d = ax + my$ certain integers x, y. This yields $d \in m\Omega$. We get $d\Omega = m\Omega$ and the minimality of m implies $m = d$. □

For every $n \in \mathbb{N}$ we can define $g(n)$ as the minimal positive generator of $n\Omega$. Put $\mathcal{A} = \{g(n); n \in \mathbb{N}\}$.

The set $r + m\Omega$ is closed and so from (2) we see that also open, so called *clopen*.

It is easy to check that the set $\mathcal{A}$ is CD - set.

Let $\{a_n\}$ be the sequence of positive integers given in (1). Clearly

$$\bigcap_{n=1}^{\infty} a_n\Omega = \{0\},$$

this yields

$$\bigcap_{m \in \mathcal{A}} m\Omega = \{0\}. \tag{3}$$

And so we ge that the set $\mathcal{A}$ is infinite. Since $m\Omega$ is open for $m \in \mathcal{A}$ equality (3) implies that for each sequence $\{\alpha_n\}$ there holds

$$\alpha_n \to 0 \Longleftrightarrow \forall m \in \mathcal{A} \exists n_0; n \geq n_0 \Longrightarrow m|\alpha_n.$$

If we define the congruence by the natural manner: $\alpha \equiv \beta \pmod{\gamma}$ if and only if γ divides $\alpha - \beta$, for $\alpha, \beta, \gamma \in \Omega$, then there holds :

$$\alpha_n \to \beta \Longleftrightarrow \forall m \in \mathcal{A} \exists n_0; n \geq n_0 \Longrightarrow \alpha_n \equiv \beta \pmod{m}.$$

Thus the convergence can be metrised by the generalized polyadic norm : Let $\mathcal{A} = \{m_n, n = 1, 2, ...\}$ and $M_n = [m_1, ..., m_n], n = 1, 2, ...$ then

$$||\alpha||_{\mathcal{A}} = \sum_{n=1}^{\infty} \frac{1 - \mathcal{X}_{M_n\Omega}(\alpha)}{2^{-n}}.$$

We get

Theorem 1. The metric ρ is equivalent with the metric $\rho_{\mathcal{A}}$ where $\rho_{\mathcal{A}}(\alpha, \beta) = ||\alpha - \beta||_{\mathcal{A}}$ for $\alpha, \beta \in \Omega$.

And so every set $S \subset \Omega$ we have

$$cl(S) = \bigcap_{n=1}^{\infty} (S + M_n\Omega) \tag{3}.$$

Denote by P the Haar probability measure defined on $(\Omega, +)$. For $m \in \mathcal{A}$ the decomposition (2) is disjoint and so $P(r + m\Omega) = \frac{1}{m}$. If we define the submeasure ν^* on the system of subsets of $\mathbb{N}$ as $\nu^*(S) = P(cl(S))$ we get from (3) and upper semicontinuity of measure that for each S

$$\nu^*(S) = \lim_{n\to\infty} \frac{R(S : M_n)}{M_n}$$

where $R(S : M_n)$ the number of elements of S incongruent modulo M_n.

Theorem 2. Let $\alpha, \beta \in \Omega$. There exist $\alpha_1, \beta_1 \in \Omega$ that the element $\delta = \alpha_1\alpha + \beta_1\beta$ divides α and β.

Proof. Let $\{a_n\}, \{b_n\}$ be the sequences of positive integers that $a_n \to \alpha, b_n \to \beta$. Let d_n the greatest common divisor of $a_n, b_n, n = 1, 2, ...$ Then $d_n = v_n a_n + u_n b_n$ for some u_n, v_n - integers. The compactness of Ω provides that $u_{k_n} \to \alpha_1$ and $v_{k_n} \to \beta_1$ for suitable increasing sequence $\{k_n\}$. Put $\delta = \alpha_1\alpha + \beta_1\beta$. We see that $d_{k_n} \to \delta$. For $n = 1, 2, ...$ we have $a_{k_n} = c_n d_{k_n}$. Since $\{c_n\}$ contains a convergent subsequence we get that δ divides α. Analogously can be derived that δ divides β. □

The element δ from Theorem 2 will be called the *greatest common divisor* of α, β and we shall write $\delta \sim (\alpha, \beta)$.

Corollary 1. If $p \in \mathcal{A}$ is a prime then for every $\alpha \in \Omega$ there holds p divides α or $(\alpha, p) \sim 1$.

Proof. If p does not divide α then $\alpha \in \Omega \setminus p\Omega$. Consider a sequence of positive integers $\{a_n\}$ which converges to α. The set $\Omega \setminus p\Omega$ is open, thus we can suppose that $(a_n, p) = 1$. This yields $\ell_n a_n + s_n p = 1$ for suitable integers ℓ_n, s_n. Since Ω is compact space there exists an increasing sequence $\{k_n\}$ that $\ell_{k_n} \to \lambda, s_{k_n} \to \sigma$. And so $\lambda\alpha + \sigma p = 1$. □

Analogously can be proved

Corollary 2. An element $\alpha \in \Omega$ is invertible if and only if $(\alpha, p) \sim 1$ for every prime $p \in \mathcal{A}$.

Lemma 2. Each closed ideal in Ω is principal ideal.

Proof. Let $I \subset \Omega$ be closed ideal. Let $\alpha \in I$. Denote by I_α he set of all divisors of α belonging to I. The compactness of Ω yields that I_α is a closed set. From Lemma 2 we get that $\alpha, \beta \in I$ there exists $\delta \in I$ that $I_\delta \subset I_\alpha \cap I_\beta$. And so by induction can be proved that $I_\alpha, \alpha \in I$ is a centered system of closed sets. Thus its intersection in non empty, it contains some element γ. Then $I = \gamma\Omega$. □

In the sequel we denote Ω the ring of polyadic integers, thus completion of $\mathbb{N}$ with respect to norm $||\cdot||_{\mathbb{N}}$ and we suppose that a infinite CD - set A is given. The completion of $\mathbb{N}$ with respect to norm $||\cdot||_A$ we denote as Ω_A.

Lemma 2 provides that $\cap_{a\in A} a\Omega = \alpha\Omega$ for suitable $\alpha \in \Omega$. We prove

Theorem 3. The ring Ω_A is isomorphic with the factor ring $\Omega/\alpha\Omega$.

Proof. There holds

If a sequence of positive integers is Cauchy's with respect to $||\cdot||_{\mathbb{N}}$ then it is Cauchy's with respect to $||\cdot||_A$ also.

If $\{a_n\}, \{b_n\}$ are sequences of positive integer then $|||a_n - b_n|||_{\mathbb{N}} \to 0 \Longrightarrow |||a_n - b_n|||_A \to 0$.

Therefore we can define a mapping $F : \Omega \to \Omega_A$ in the following way: If $\beta \in \Omega, b_n \to \beta$ with respect $||\cdot||_{\mathbb{N}}$ then $F(\beta)$ is the limit of $\{b_n\}$ with respect to $||\cdot||_A$. Clearly F is a surjective morphism with kernel $\cap_{a\in A} a\Omega$ and the assertion follows. □

References

[N] NOVOSELOV, E. V., *Topological theory of polyadic numbers*, Trudy Tbilis. Mat. Inst. 27, 61 - 69, (1960), in russian

[N1] NOVOSELOV, E. V., *New method in the probability number theory*, Doklady akademii nauk. ser. matem. No. 2, 28, 307 - 364, (1964), in russian

[P] PASTEKA, M., *Covering densities*, Math. Slovaca 42 (1992), 593-614.

[Po] R.G. POSTNIKOV, *Introduction to analytic number theory, (in russian)*, Moscov, Nauka, 1971, english translation: Amer Math Soc, Providence RI 1981

[P-P] PAŠTÉKA, M. PORUBSKÝ, Š., *On the distribution on the sequences of integers.*, Math. Slovaca 43 (1993), 521-639

Katedra matematiky a informatiky, Pedagogicka Fakulta, Trnavska Univerzita
Priemyselna 4, TRNAVA, SLOVAKIA
e - mail: pasteka@mat.savba.sk